



CONSEIL AFRICAIN ET MALGACHE POUR L'ENSEIGNEMENT SUPÉRIEUR
01 BP 134 OUAGADOUGOU 01 (BURKINA FASO)
TEL (226) 25 36 81 46 - FAX (226) 25 36 85 73 - Email: comes@lecomes.org

COMITES CONSULTATIFS INTERAFRICAINS

Article 2

Position sur l'article : 3

Nom sur la publication : **Ballo Daouda**



CONSEIL AFRICAIN ET MALGACHE POUR L'ENSEIGNEMENT SUPÉRIEUR
01 BP 134 OUAGADOUGOU 01 (BURKINA FASO)
TEL (226) 25 36 81 46 - FAX (226) 25 36 85 73 - Email: comes@lecomes.org

COMITES CONSULTATIFS INTERAFRICAINS

PUBLICATION N°2

Titre: 3-[2-(3-Methylquinoxalin-2-yloxy) ethyl]-1,3-oxazolidin-2-one

Date : 2010

Auteurs: Caleb Anothane Ahoya, Rachid Bouhfid, **Ballo Daouda**, El Mokhtar Essassi and Lahcen El Ammari

Nom de la revue scientifique: Acta Crystallographica, Section E: Structure Reports Online

En ligne: <https://journals.iucr.org/e/index.html>

Volume: 66

Issue: 5

Page: o1050

ISSN: 1600-5368 (ancien)

ISSN: 2056-9890 (nouveau)

Maison d'édition : International Union of Crystallography

Facteur d'impact: 0,34

Indexation: Chemical Abstracts, Scopus (Elsevier)

CODEN : ACSEBH (Ancien), ACSECI (nouveau)

Sourcerecord id : 21100427192



CONSEIL AFRICAIN ET MALGACHE POUR L'ENSEIGNEMENT SUPÉRIEUR
01 BP 134 OUAGADOUGOU 01 (BURKINA FASO)
TEL (226) 25 36 81 46 - FAX (226) 25 36 85 73 - Email: comes@lecomes.org

COMITES CONSULTATIFS INTERAFRICAINS

Preuves d'indexation



CONSEIL AFRICAIN ET MALGACHE POUR L'ENSEIGNEMENT SUPÉRIEUR
01 BP 134 OUAGADOUGOU 01 (BURKINA FASO)
TEL (226) 25 36 81 46 - FAX (226) 25 36 85 73 - Email: comes@lecomes.org

COMITES CONSULTATIFS INTERAFRICAINS




Acta Crystallographica Section E

CRYSTALLOGRAPHIC COMMUNICATIONS



search IUCr Journals

home archive editors for authors for readers submit open access




Acta Crystallographica Section E: Crystallographic Communications is the IUCr's open-access structural communications journal. It provides a fast, simple and easily accessible publication mechanism for crystal structure determinations of inorganic, metal-organic and organic compounds. The electronic submission, validation, refereeing and publication facilities of the journal ensure rapid and high-quality publication of fully validated structures. The primary article category is *Research Communications*; these are peer-reviewed articles describing one or more structure determinations with appropriate discussion of the science. Note that *Data Reports* are now published in *IUCrData*.

More about the journal

Editorial board
ISSN: 2056-9890

Special issues



Weak interactions

Weak interactions in crystals:
an integrated approach



CONSEIL AFRICAIN ET MALGACHE POUR L'ENSEIGNEMENT SUPÉRIEUR
01 BP 134 OUAGADOUGOU 01 (BURKINA FASO)
TEL (226) 25 36 81 46 - FAX (226) 25 36 85 73 - Email: comes@lecomes.org

COMITES CONSULTATIFS INTERAFRICAINS



Acta Crystallographica Section E

CRYSTALLOGRAPHIC COMMUNICATIONS

[home](#) [archive](#) [editors](#) [for authors](#) [for readers](#) [submit](#) [open access](#)

AIMS AND SCOPE

Acta Crystallographica Section E: Crystallographic Communications is the IUCr's open-access structural communications journal. It provides a fast, simple and easily accessible publication mechanism for crystal structure determinations of inorganic, metal-organic and organic compounds. The electronic submission, validation, refereeing and publication facilities of the journal ensure rapid and high-quality publication of fully validated structures. The primary article category is *Research Communications*; these are peer-reviewed articles describing one or more structure determinations with appropriate discussion of the science. Note that *Data Reports* are now published in *IUCrData*.

JOURNAL IDENTIFIERS

Title: *Acta Crystallographica Section E*
Subtitle: *Crystallographic Communications*
Abbreviated title: *Acta Cryst. E*
ISSN: 2056-9890
CODEN: ACSECI
DOI: <https://doi.org/10.1107/S20569890>

ABSTRACTING DETAILS

Acta Crystallographica Section E is covered by:

- Ceramic Abstracts
- Chemical Abstracts
- Cambridge Structural Database
- Emerging Sources Citation Index
- Inorganic Crystal Structure Database
- INSPEC
- Medline
- Metals Abstracts/METADEx
- Scopus



CONSEIL AFRICAIN ET MALGACHE POUR L'ENSEIGNEMENT SUPÉRIEUR
01 BP 134 OUAGADOUGOU 01 (BURKINA FASO)
TEL (226) 25 36 81 46 - FAX (226) 25 36 85 73 - Email: comes@lecomes.org

COMITES CONSULTATIFS INTERAFRICAINS



CASSISM
A CAS SOLUTION



CAS Source Index (CASSI) Search Tool

[Search](#) | [About](#) | [Contact Us](#) | [Help](#) | [CAS](#) | [American Chemical Society](#)

CAS Source Index (CASSI) Search Result

Displaying Record for Publication: [Acta Crystallographica, Section E: Structure Reports Online](#)

Entry Type	Changed Title Serial
Title	Acta Crystallographica, Section E: Structure Reports Online
Abbreviated Title	Acta Crystallogr., Sect. E: Struct. Rep. Online
CODEN	ACSEBH
ISSN	1600-5368
Language of Text	English
History	vE57 n1 2001-vE70 n12 Dec. 2014
Publication Notes	Avail. from Internet at URL: http://journals.iucr.org/e/journalhomepage.html
Publisher Name	International Union of Crystallography
Successor Title Note	Changed to
Successor Title(s)	Acta Crystallographica, Section E: Crystallographic Communications
Alternate Title(s)	Structure Reports Online
Abbreviated Alternate Title(s)	Struct. Rep. Online

Disclaimer

[Search](#) | [About](#) | [Contact Us](#) | [Help](#) | [CAS](#) | [American Chemical Society](#)



CAS[®]
A DIVISION OF THE
AMERICAN CHEMICAL SOCIETY

Copyright © 2020 American Chemical Society All Rights Reserved



CONSEIL AFRICAIN ET MALGACHE POUR L'ENSEIGNEMENT SUPÉRIEUR
01 BP 134 OUAGADOUGOU 01 (BURKINA FASO)
TEL (226) 25 36 81 46 - FAX (226) 25 36 85 73 - Email: comes@lecomes.org

COMITES CONSULTATIFS INTERAFRICAINS



CASSISM
A CAS SOLUTION



CAS Source Index (CASSI) Search Tool

[Search](#) | [About](#) | [Contact Us](#) | [Help](#) | [CAS](#) | [American Chemical Society](#)

CAS Source Index (CASSI) Search Result

Displaying Record for Publication: [Acta Crystallographica, Section E: Crystallographic Communications](#)

Entry Type	Active Serial
Title	Acta Crystallographica, Section E: Crystallographic Communications
Abbreviated Title	Acta Crystallogr., Sect. E: Crystallogr. Commun.
CODEN	ACSECI
ISSN	2056-9898
Former Title Note(s)	Formerly
Former Title(s)	Acta Crystallographica, Section E: Structure Reports Online
Language of Text	English
Summaries In	English
History	vE71 n1 Jan. 2015+
Publication Notes	From 2016 Data Reports pub. in IUCrData, which see; Avail. from Internet at URL: http://journals.iucr.org/e/
Publisher Name	International Union of Crystallography

Disclaimer

[Search](#) | [About](#) | [Contact Us](#) | [Help](#) | [CAS](#) | [American Chemical Society](#)



CAS[®]
A DIVISION OF THE
AMERICAN CHEMICAL SOCIETY

Copyright © 2020 American Chemical Society All Rights Reserved



CONSEIL AFRICAIN ET MALGACHE POUR L'ENSEIGNEMENT SUPÉRIEUR
01 BP 134 OUAGADOUGOU 01 (BURKINA FASO)
TEL (226) 25 36 81 46 - FAX (226) 25 36 85 73 - Email: comes@lecomes.org

COMITES CONSULTATIFS INTERAFRICAINS

1	Source Title (CSA excl.) (Medline-sourced journals are indicated in Green). Including Conference Proceedings available in the scopus.com Source Browse list	Print-ISSN	E-ISSN	Coverage
230	Acta Crystallographica Section E: Structure Reports	16005368		2003-ongoing

1	Sourcerecord id	Source Title (Medline-sourced journals are indicated in Green) Titles indicated in bold red do not meet the Scopus quality criteria anymore and therefore Scopus discontinued the forward capturing	Print-ISSN	E-ISSN	Active or Inactive	Coverage	Article language in source (three-letter ISO language)
334	21100427192	Acta Crystallographica Section E: Crystallographic Communications		20569890	Active	2015-ongoing	ENG



CONSEIL AFRICAIN ET MALGACHE POUR L'ENSEIGNEMENT SUPÉRIEUR
01 BP 134 OUAGADOUGOU 01 (BURKINA FASO)
TEL (226) 25 36 81 46 - FAX (226) 25 36 85 73 - Email: comes@lecomes.org

COMITES CONSULTATIFS INTERAFRICAINS


Article : 2



CONSEIL AFRICAIN ET MALGACHE POUR L'ENSEIGNEMENT SUPÉRIEUR
01 BP 134 OUAGADOUGOU 01 (BURKINA FASO)
TEL (226) 25 36 81 46 - FAX (226) 25 36 85 73 - Email: comes@lecomes.org

COMITES CONSULTATIFS INTERAFRICAINS

Acta Cryst. (2010). E66, o1050
<https://doi.org/10.1107/S1600536810012687>




3-[2-(3-Methylquinoxalin-2-yloxy)ethyl]-1,3-oxazolidin-2-one

C. A. Ahoya, R. Bouhfid, B. Daouda, E. M. Essassi and L. El Ammari

[Read article](#)

Acta Cryst. (2010). E66, o1051-o1052
<https://doi.org/10.1107/S1600536810011918>



Ethyl 1-tert-butyl-2-(4-hydroxy-3-methoxyphenyl)-1H-benzimidazole-5-carboxylate

N. Arumugam, A. S. Abdul Rahim, H. Osman, M. M. Rosli and H.-K. Fun

[Read article](#)

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

3-[2-(3-Methylquinoxalin-2-yloxy)ethyl]-1,3-oxazolidin-2-one

Caleb Anothane Ahoya,^a Rachid Bouhfid,^b Ballo Daouda,^a El Mokhtar Essassi^{a*} and Lahcen El Ammari^c

^aLaboratoire de Chimie Organique Hétérocyclique, Pôle de Compétences, Pharmacochimie, Avenue Ibn Battouta, BP 1014, Faculté des Sciences, Université Mohammed V-Agdal, Rabat, Morocco, ^bInstitute of Nanomaterial and Nanotechnology, Avenue Armée Royale, Rabat, Morocco, and ^cLaboratoire de Chimie du Solide Appliquée, Faculté des Sciences, Université Mohammed V-Agdal, Avenue Ibn Battouta, BP 1014, Rabat, Morocco

Correspondence e-mail: emessassi@yahoo.fr


Received 27 March 2010; accepted 5 April 2010

Key indicators: single-crystal X-ray study; $T = 296\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; R factor = 0.040; wR factor = 0.121; data-to-parameter ratio = 15.3.

Two isomers were isolated during the reaction between 3-methylquinoxalin-2-one and bis(2-chloroethyl)amine hydrochloride. The crystal structure of one isomer has already been reported [Caleb, Bouhfid, Essassi & El Ammari (2009). *Acta Cryst. E* **65**, o2024–o2025], while that of the second isomer is the subject of this work. The title compound, $\text{C}_{14}\text{H}_{15}\text{N}_3\text{O}_3$, has a new structure containing oxazolidine and quinoxaline rings linked by an ethoxy group. The main difference between the two isomers is the position of the oxazolidine group with respect to the quinoxaline system. The dihedral angle between the fused planar rings and the oxazolidin-2-one ring is $41.63(8)^\circ$ in the title molecule.

Related literature

For the biological activity of 3-[2-(3-methyl-1,2-dihydroquinoxalin-2-yloxy)ethyl]oxazolidin-2-one, see: Madhusudhan *et al.* (2004); Soad *et al.* (2006); Sriharsha & Shashikanth (2006); Menoret *et al.* (2009); Wilhelmsson *et al.* (2008). For the structure of the isomer of the title compound, see: Caleb *et al.* (2009). For related structures, see: Doubia *et al.* (2007); Mamedov *et al.* (2007); Aschwanden *et al.* (1976)



* Present address: Institute of Nanomaterial and Nanotechnology, Avenue de l'Armée Royale, Rabat, Morocco.

Experimental

Crystal data

$\text{C}_{14}\text{H}_{15}\text{N}_3\text{O}_3$
 $M_r = 273.29$
 Triclinic, $P\bar{1}$
 $a = 6.9936(3)\text{ \AA}$
 $b = 7.6916(3)\text{ \AA}$
 $c = 13.3709(6)\text{ \AA}$
 $\alpha = 86.649(2)^\circ$
 $\beta = 77.044(2)^\circ$
 $\gamma = 71.141(2)^\circ$
 $V = 663.23(5)\text{ \AA}^3$
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 0.10\text{ mm}^{-1}$
 $T = 296\text{ K}$
 $0.41 \times 0.33 \times 0.20\text{ mm}$

Data collection

Bruker X8 APEXII CCD area-detector diffractometer
 15358 measured reflections
 3030 independent reflections
 2358 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.023$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.040$
 $wR(F^2) = 0.121$
 $S = 1.06$
 3030 reflections
 198 parameters
 H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.21\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.17\text{ e \AA}^{-3}$

Data collection: APEX2 (Bruker, 2005); cell refinement: APEX2; data reduction: APEX2; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 for Windows (Farrugia, 1997) and PLATON (Spek, 2009); software used to prepare material for publication: WinGX (Farrugia, 1999).

The authors thank the Unit of Support for Technical and Scientific Research (UATRS, CNRST) for the X-ray measurements.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: DN2552).

References

- Aschwanden, W., Kyburz, E. & Schonholzer, P. (1976). *Helv. Chim. Acta*, **59**, 1245–1252.
 Bruker (2005). APEX2. Bruker AXS Inc., Madison, Wisconsin, USA.
 Caleb, A. A., Bouhfid, R., Essassi, E. M. & El Ammari, L. (2009). *Acta Cryst. E* **65**, o2024–o2025.
 Doubia, M. L., Bouhfid, R., Ahabchane, N. H., Essassi, E. M. & El Ammari, L. (2007). *Acta Cryst. E* **63**, o3305.
 Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
 Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
 Madhusudhan, G., Om Reddy, G., Ramanathan, J. & Dubey, P. K. (2004). *Indian J. Chem. Sect. B*, **43**, 957–963.
 Mamedov, V. A., Khafizova, E. A., Beschastnova, T. N., Zhukova, N. A., Gubaidullin, A. T., Rizvanov, I. Kh., Berdnikov, E. A. & Litvinov, I. A. (2007). *Izv. Akad. Nauk SSSR Ser. Khim. (Russ.) (Russ. Chem. Bull.)*, **5**, 1047–1048.
 Menoret, A., Mcaleer, J. P., Ngoi, S. M., Swagatam, R., Eddy, N. A., Fenteany, G., Lee, S. J., Rossi, R. J., Mukherji, B., Allen, D. L., Chakraborty, N. G. & Vella, A. T. (2009). *J. Immunol.* **183**, 7489–7496.
 Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
 Soad, A. M., El-Hawash, S. A., Abeer, E. & Wahab, A. (2006). *Arch. Pharm. (Weinheim)*, **339**, 437–447.
 Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
 Sriharsha, S. N. & Shashikanth, S. J. (2006). *Heterocycl. Commun.* **12**, 213–218.
 Wilhelmsson, L. M., Kingi, N. & Bergman, J. (2008). *J. Med. Chem.* **51**, 7744–7750.

supporting information

Acta Cryst. (2010). E66, o1050 [doi:10.1107/S1600536810012687]

3-[2-(3-Methylquinoxalin-2-yloxy)ethyl]-1,3-oxazolidin-2-one

Caleb Anothane Ahoya, Rachid Bouhfid, Ballo Daouda, El Mokhtar Essassi and Lahcen El Ammari

S1. Comment

Oxazolidin-2-ones and quinoxalines are subjects of numerous articles in scientific journals concerning the development of new molecules as drug candidates such as antibacterials (Madhusudhan *et al.* 2004); (Sriharsha & Shashikanth, 2006), anti-viral (Wilhelmsson, *et al.* 2008), anti-tumor (Soad *et al.* 2006), and anti-inflammatory (Menoret *et al.* 2009). Our investigation is intended to increase the biological activity of such molecules. During the synthesis, two isomers were isolated, and the structure of isomer 1 has already been published (Caleb *et al.* 2009) while that of isomer 2 is the subject of the present work.


The structure of the 3-(2-(3-methyl-1,2-dihydro-quinoxalin-2-yloxy)ethoxy) oxazolidin-2-one molecule is also built up from two fused six-membered rings linked to a five-membered ring (oxazolidin-2-one) by an ethoxy group, as shown in Fig.1. It would be interesting to compare the crystal structures of both isomers of this compound (scheme 1). Actually, the geometric parameters (bond lengths and angles) of the two isomers are very similar to those observed in other heterocyclic structures (Aschwenden *et al.*, 1976; Doubia *et al.*, 2007; Mamedov *et al.*, 2007). However, the main difference between the two isomers is the position of the oxazolidine group with respect to the quinoxalin. Moreover, the dihedral angle between the fused six-membered rings and the five cycles measuring 20.04 (9)° in the isomer 1 instead of 41.63 (8)° in the isomer 2.

S2. Experimental


In a 100 ml flask, is reacted 0.0125 moles of quinoxalin-2-one with 2.66 moles of dichloroethylamine hydrochloride in 40 ml of dimethyl formamide in presence of 2.87 moles of potassium carbonate and a few milligrams of tetran-butyl ammonium bromide. The mixture was brought to reflux in a sand bath, magnetic stirring and the reaction progress was monitored by thin layer chromatography. After evaporation of solvent under reduced pressure, the residue obtained is chromatographed on silica column (hexane / acetate: 4 / 6). Thus we have isolated two compounds. Recrystallization occurred in the same eluent. This compound was obtained in 38% and his melting point is 169°C.

S3. Refinement

H atoms were located in a difference map and treated as riding with C—H = 0.96 Å for methyl groups and C—H = 0.93 Å for all other hydrogens with $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{aromatic, methine})$ or $U_{\text{iso}}(\text{H}) = 1.5 U_{\text{eq}}(\text{methyl})$. All other H atoms were located from difference Fourier maps and refined without any distance restraints.

**Figure 1**

Molecular structure of the title compound with the atom-labelling scheme. Displacement ellipsoids are drawn at the 50% probability level. H atoms are represented as small circles.

**Figure 2**

The structures of the two isomers.

3-[2-(3-Methylquinoxalin-2-yloxy)ethyl]-1,3-oxazolidin-2-one

Crystal data

$C_{14}H_{15}N_3O_3$

$M_r = 273.29$

Triclinic, $P\bar{1}$

Hall symbol: $-P\ 1$

$a = 6.9936\ (3)\ \text{\AA}$

$b = 7.6916\ (3)\ \text{\AA}$

$c = 13.3709\ (6)\ \text{\AA}$

$\alpha = 86.649\ (2)^\circ$

$\beta = 77.044\ (2)^\circ$

$\gamma = 71.141\ (2)^\circ$

$V = 663.23 (5) \text{ \AA}^3$
 $Z = 2$
 $F(000) = 288$
 $D_x = 1.368 \text{ Mg m}^{-3}$
 Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
 Cell parameters from 15358 reflections

$\theta = 2.8\text{--}27.5^\circ$
 $\mu = 0.10 \text{ mm}^{-1}$
 $T = 296 \text{ K}$
 Prism, colourless
 $0.41 \times 0.33 \times 0.20 \text{ mm}$

Data collection

Bruker X8 APEXII CCD area-detector
 diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 φ and ω scans
 15358 measured reflections
 3030 independent reflections

2358 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.023$
 $\theta_{\text{max}} = 27.5^\circ$, $\theta_{\text{min}} = 2.8^\circ$
 $h = -9 \rightarrow 9$
 $k = -9 \rightarrow 9$
 $l = -17 \rightarrow 17$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.040$
 $wR(F^2) = 0.121$
 $S = 1.06$
 3030 reflections
 198 parameters
 0 restraints
 Primary atom site location: structure-invariant
 direct methods

Secondary atom site location: difference Fourier
 map
 Hydrogen site location: inferred from
 neighbouring sites
 H atoms treated by a mixture of independent
 and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0643P)^2 + 0.0716P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} < 0.001$
 $\Delta\rho_{\text{max}} = 0.21 \text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.17 \text{ e \AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ($\text{\AA}^2$)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.31122 (17)	0.72867 (15)	0.95763 (8)	0.0378 (3)
C2	0.08813 (17)	0.80823 (15)	1.11146 (8)	0.0382 (3)
C3	-0.11007 (19)	0.83829 (18)	1.17492 (10)	0.0470 (3)
C4	-0.1442 (2)	0.8837 (2)	1.27638 (10)	0.0531 (3)
C5	0.0143 (2)	0.90272 (19)	1.31801 (10)	0.0540 (3)
C6	0.2068 (2)	0.87655 (18)	1.25750 (10)	0.0503 (3)
C7	0.24799 (18)	0.82807 (16)	1.15335 (9)	0.0403 (3)
C8	0.47691 (17)	0.75240 (16)	0.99773 (9)	0.0412 (3)
C9	0.68526 (19)	0.7217 (2)	0.92895 (11)	0.0546 (3)
H9A	0.7371	0.5987	0.9015	0.082*
H9B	0.7784	0.7393	0.9674	0.082*

H9C	0.6743	0.8075	0.8737	0.082*
C10	0.21267 (19)	0.63937 (19)	0.81444 (9)	0.0465 (3)
H10A	0.1105	0.7551	0.8041	0.056*
H10B	0.1429	0.5643	0.8593	0.056*
C11	0.3226 (2)	0.54186 (19)	0.71328 (10)	0.0523 (3)
H11A	0.4218	0.4258	0.7258	0.063*
H11B	0.2221	0.5146	0.6826	0.063*
C12	0.3726 (3)	0.7066 (2)	0.55269 (11)	0.0625 (4)
C13	0.6852 (3)	0.7454 (3)	0.54185 (14)	0.0820 (5)
H13A	0.6957	0.8634	0.5570	0.098*
H13B	0.8151	0.6739	0.4986	0.098*
C14	0.6363 (3)	0.6450 (3)	0.63974 (11)	0.0682 (4)
H14A	0.7319	0.5209	0.6375	0.082*
H14B	0.6394	0.7098	0.6991	0.082*
N1	0.12438 (14)	0.75658 (13)	1.01023 (7)	0.0406 (2)
N2	0.44386 (15)	0.80020 (15)	1.09343 (8)	0.0462 (3)
N3	0.43028 (18)	0.64507 (16)	0.64066 (8)	0.0534 (3)
O1	0.36793 (12)	0.67107 (12)	0.85906 (6)	0.0460 (2)
O2	0.2172 (2)	0.70974 (18)	0.52691 (9)	0.0879 (4)
O3	0.5186 (2)	0.76983 (16)	0.49204 (8)	0.0829 (4)
H3	−0.220 (2)	0.8300 (19)	1.1449 (11)	0.052 (4)*
H4	−0.280 (2)	0.908 (2)	1.3175 (13)	0.068 (4)*
H5	−0.009 (2)	0.934 (2)	1.3883 (14)	0.064 (4)*
H6	0.322 (3)	0.889 (2)	1.2864 (13)	0.071 (4)*

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0389 (6)	0.0433 (6)	0.0313 (6)	−0.0140 (5)	−0.0066 (4)	0.0011 (4)
C2	0.0413 (6)	0.0410 (6)	0.0317 (6)	−0.0135 (5)	−0.0066 (4)	0.0011 (4)
C3	0.0439 (7)	0.0559 (7)	0.0406 (7)	−0.0189 (6)	−0.0031 (5)	−0.0016 (5)
C4	0.0526 (7)	0.0592 (8)	0.0396 (7)	−0.0161 (6)	0.0039 (6)	−0.0019 (6)
C5	0.0656 (8)	0.0568 (8)	0.0317 (7)	−0.0106 (6)	−0.0064 (6)	−0.0048 (5)
C6	0.0545 (8)	0.0564 (8)	0.0388 (7)	−0.0113 (6)	−0.0162 (6)	−0.0043 (5)
C7	0.0415 (6)	0.0427 (6)	0.0354 (6)	−0.0106 (5)	−0.0097 (5)	−0.0003 (5)
C8	0.0366 (6)	0.0476 (6)	0.0391 (6)	−0.0127 (5)	−0.0083 (5)	−0.0005 (5)
C9	0.0394 (6)	0.0745 (9)	0.0505 (8)	−0.0210 (6)	−0.0050 (5)	−0.0066 (6)
C10	0.0446 (6)	0.0629 (8)	0.0351 (6)	−0.0211 (6)	−0.0079 (5)	−0.0038 (5)
C11	0.0608 (8)	0.0600 (8)	0.0381 (7)	−0.0217 (6)	−0.0093 (6)	−0.0066 (5)
C12	0.0842 (11)	0.0549 (8)	0.0358 (7)	−0.0028 (8)	−0.0139 (7)	−0.0097 (6)
C13	0.0972 (13)	0.0874 (12)	0.0545 (10)	−0.0363 (10)	0.0076 (9)	0.0021 (8)
C14	0.0723 (10)	0.0929 (11)	0.0440 (8)	−0.0373 (9)	−0.0061 (7)	0.0036 (7)
N1	0.0385 (5)	0.0515 (6)	0.0331 (5)	−0.0169 (4)	−0.0062 (4)	−0.0017 (4)
N2	0.0400 (5)	0.0571 (6)	0.0425 (6)	−0.0140 (5)	−0.0120 (4)	−0.0036 (5)
N3	0.0602 (7)	0.0644 (7)	0.0316 (5)	−0.0157 (5)	−0.0071 (5)	−0.0024 (5)
O1	0.0409 (4)	0.0667 (6)	0.0316 (4)	−0.0203 (4)	−0.0035 (3)	−0.0063 (4)
O2	0.1019 (9)	0.0914 (9)	0.0633 (8)	−0.0034 (7)	−0.0421 (7)	−0.0077 (6)
O3	0.1211 (10)	0.0797 (8)	0.0399 (6)	−0.0288 (7)	−0.0081 (6)	0.0096 (5)

Geometric parameters (Å, °)

C1—N1	1.2932 (14)	C9—H9C	0.9600
C1—O1	1.3457 (13)	C10—O1	1.4398 (13)
C1—C8	1.4458 (15)	C10—C11	1.5041 (18)
C2—N1	1.3777 (14)	C10—H10A	0.9700
C2—C3	1.4083 (16)	C10—H10B	0.9700
C2—C7	1.4099 (15)	C11—N3	1.4523 (17)
C3—C4	1.3695 (18)	C11—H11A	0.9700
C3—H3	0.966 (14)	C11—H11B	0.9700
C4—C5	1.397 (2)	C12—O2	1.2046 (19)
C4—H4	0.949 (16)	C12—N3	1.3394 (19)
C5—C6	1.3646 (19)	C12—O3	1.357 (2)
C5—H5	0.948 (17)	C13—O3	1.424 (2)
C6—C7	1.4045 (17)	C13—C14	1.510 (2)
C6—H6	0.999 (17)	C13—H13A	0.9700
C7—N2	1.3793 (15)	C13—H13B	0.9700
C8—N2	1.3013 (15)	C14—N3	1.4379 (19)
C8—C9	1.4923 (16)	C14—H14A	0.9700
C9—H9A	0.9600	C14—H14B	0.9700
C9—H9B	0.9600		
N1—C1—O1	121.60 (10)	C11—C10—H10A	110.3
N1—C1—C8	124.34 (10)	O1—C10—H10B	110.3
O1—C1—C8	114.06 (9)	C11—C10—H10B	110.3
N1—C2—C3	119.79 (10)	H10A—C10—H10B	108.6
N1—C2—C7	120.95 (10)	N3—C11—C10	114.20 (11)
C3—C2—C7	119.25 (11)	N3—C11—H11A	108.7
C4—C3—C2	119.75 (12)	C10—C11—H11A	108.7
C4—C3—H3	121.5 (9)	N3—C11—H11B	108.7
C2—C3—H3	118.7 (8)	C10—C11—H11B	108.7
C3—C4—C5	121.01 (12)	H11A—C11—H11B	107.6
C3—C4—H4	118.6 (10)	O2—C12—N3	127.96 (16)
C5—C4—H4	120.3 (10)	O2—C12—O3	122.31 (14)
C6—C5—C4	120.14 (12)	N3—C12—O3	109.73 (14)
C6—C5—H5	118.9 (9)	O3—C13—C14	106.05 (14)
C4—C5—H5	120.9 (9)	O3—C13—H13A	110.5
C5—C6—C7	120.42 (12)	C14—C13—H13A	110.5
C5—C6—H6	120.9 (10)	O3—C13—H13B	110.5
C7—C6—H6	118.6 (10)	C14—C13—H13B	110.5
N2—C7—C6	119.74 (10)	H13A—C13—H13B	108.7
N2—C7—C2	120.85 (10)	N3—C14—C13	101.80 (14)
C6—C7—C2	119.41 (11)	N3—C14—H14A	111.4
N2—C8—C1	119.95 (10)	C13—C14—H14A	111.4
N2—C8—C9	120.35 (10)	N3—C14—H14B	111.4
C1—C8—C9	119.70 (11)	C13—C14—H14B	111.4
C8—C9—H9A	109.5	H14A—C14—H14B	109.3
C8—C9—H9B	109.5	C1—N1—C2	115.96 (9)

H9A—C9—H9B	109.5	C8—N2—C7	117.92 (10)
C8—C9—H9C	109.5	C12—N3—C14	112.13 (13)
H9A—C9—H9C	109.5	C12—N3—C11	122.09 (13)
H9B—C9—H9C	109.5	C14—N3—C11	123.42 (12)
O1—C10—C11	106.91 (10)	C1—O1—C10	117.34 (9)
O1—C10—H10A	110.3	C12—O3—C13	109.59 (12)
